LIBRARY
FISH & WILDLIFE SERVICE
PACIFIC OCEANIC FISHERY
INVESTIGATIONS

BULLETIN OF THE SCRIPPS INSTITUTION OF OCEANOGRAPHY

TECHNICAL SERIES

Volume III

1931-1935

EDITORS
GEORGE F. McEWEN
ERIK G. MOBERG
FRANCIS B. SUMNER
T. WAYLAND VAUGHAN

1904 60

UNIVERSITY OF CALIFORNIA PRESS
BERKELEY, CALIFORNIA
1937

UNIVERSITY OF CALIFORNIA PRESS
BERKELEY, CALIFORNIA

CAMBRIDGE UNIVERSITY PRESS
LONDON, ENGLAND

PRINTED IN THE UNITED STATES OF AMERICA

INDEX

Acartia clausii, 33
tonsa, 33
Acknowledgments, 4, 33, 117, 225, 232, 356
Actinomyces in sea water, 292
Activity coefficient: of bicarbonate and carbonate ions in sea water, 264 ff.; of calcium ion in sea water, 272. *See also* Sea water
Agassiz, Alexander, investigations of, on sea-water deposits, 4
Agassiz, Louis, work of, 4
Aglaophenia struthionides, 51
Air masses: transitional character of, 312; characteristic properties of, 312, 314; "differential" definition of, 314; stability of, 315, 319; transitional polar continental, 316; of United States compared with Europe, 320; interaction of, 320 ff.; stable, of mountainous regions, 322; frontal movements of, 323 ff.; convective instability of, 327; effect of mountain ranges on, 329, 330; real vertical structure of, 330
Air Masses of the North Pacific, The, 311–354
Alaska, cyclones of, 322.
Aletes squamigerus, 63
Aleutian Islands, 319, 322, 323, 324, 328
Algae: metabolic activities of, 47; time of appearance of, 49; bound in tubular nests by annelids, 52
Allen, Esther C. *et al.*, **The Buffer Mechanism of Sea Water,** 231–278
Alternation of generations, dimorphic tests the result of, 396
Amaroucium californicum, 71
Ameniya, I., cited on sex change in the Japanese oyster, 136
Ammonia: production of, by bacteria, 289; from peptone, 290; from urea, 290
Amoebula: spherical, 359, 363; zygotic, 359, 369, 402; cytoplasm of, 369; postzygotic, 396; proloculum surrounded by cytoplasm of, 396
Andrews, E. A., cited on development of reproductive organs, 206
Annelids: attachment of, to wood and cement, 42; time of appearance of, 52; habits of, 52; algae bound into tubular nests by, 52
Anodonta, 130
Aragonite, natural, $CaSO_4$ content of, 152
Ascidia californica, 48, 71
Asiatic continent, movement of cold occluded fronts off the, 321; interaction of air masses along the frontal zones of the, 322
Atherinopsis californicus, 89; egg mass of, 89, 92, 93; larvae of, 91; variation of color in larvae of, 91, 93
Atlantic air, polar, long free path over ocean of, 320
Atlantic, Icelandic cyclone of, 323

Bacilli found in sea water, 292
Bacteria, marine: sampling technic for, 279; colonies of, 282; enumeration of, 282; heat-sensitivity of, 282; incubation temperature for, 283; distribution of, 284–287; effect of radiation and sunlight on, 286; effect of tide on distribution of, 286; seasonal variation of, 287; in muds, 288; aerobic, 289; anaerobic, 289–291; physiological types of, 289–292; denitrifying, 290; chromogenesis of, 291; motility of, 291; morphology of, 292
Bacteriological water sampler: difficulty in producing an ideal, 191, 192, 279–281; various types of, 192; sterilization of, 192; attachment for Ekman bottle, 193 ff.; sampling operation, 197, 198, 199; danger of contamination of, 199; effect of depth on, 280
Bacteriological Water Sampler, 191–200, 280
Bailey, J. W., work on sea-bottom deposits by, 4
Balanus tintinnabulum californicus, 41, 47, 63
Barnhart, Percy S., **Notes on the Habits, Eggs, and Young of Some Fishes of Southern California,** 87–99
Bassler, R. S., identification of bryozoa by, 53
Batavia, Java, decrease in equivalent-potential temperature at, 315
Beef kidney: stability of catalase in, extract, 300; comparison between mussel catalase and that of, 306, 309; temperature at which maximum catalase activity is obtained from, 307, 308, 309

Bering Sea: a polar air source, 312, 314; wind currents of, 322, 327, 334; equivalent-potential temperature in, 327
Berkefeld filters, materials removed from sea water by, 174
Blue Muds, 5, 20; color of, 6; organic constituents of, 6; minerals observed in, 6; location of, 6, 27; sand grades of, differ from those of Red Clays, 12
Bolivina subadvena var. spissa, 228
argentea, 288
Boron, rôle of, in buffer mechanism of sea water, 260 ff.
Boston, upper air soundings made at, 317
Botula (Adula) diegensis, 62
Branchiomma disparoculatum, 52
Bryozoa, growth of, 53
Buch *et al.*, investigations on CO_2 of sea water by, 232, 235
Buch, K., cited on: pH and CO_2 of sea water, 232; factors involved in buffer mechanism, 233, 234
Buchanan, J. Y., hydrometer described by, 101
Buffer Mechanism of Sea Water, The, 231–278
Buffer mechanism of sea water: quantitative formulation of, 235; interconnected factors of, 235; divalent ions of, 235; mass law equations of, 236 ff., 244 ff.; rôle of boron in, 260 ff. *See also* Mass equations; Sea water
Bugula, 42, 47
eburnea, 47, 63
flabellata, 53
neritina, 47, 53, 54
Bulimina, 403
rostrata, 228 *passim*
Buliminella elegantissima, 227
Burkenroad, M. D. cited on sexual phases in Ostrea virginica, 137
Burrowing mechanism of Dendrostoma zostericola, 210
Byers, Horace R., **The Air Masses of the North Pacific,** 311–354

Calcareous Mud, 5, 15; color of, 9; organisms found in, 9; minerals observed in, 9; criteria for recognition of, 9; West Indian deposits of, 9; patchy distribution of, 9
Calcium carbonate: determination of, 147; solubility product constant of, 150, 153; conditions of oversaturation with, 151; four different forms of, 152; deposition of, caused by death of organisms, 153
Calcium Equilibrium in Sea Water, 145–190
Calcium equilibrium in sea water: saturation and undersaturation, 145, 151, 158, 164, 180, 185; relationships of, 145, 155; physiochemical behavior of, 146; liquid phase of, 147; limit of solubility in, 147; kinetic theory consideration of, 147, 151; limit for molar concentration in, 147; application of ionic concept to, 149; in gaseous and liquid phases, 152, 153; effect of biological activity on, 153–154, 162, 174, 179; influence of solid phase on, 152–153, 161, 186; effect of reduction of CO_2 content on, 162–163, 168, 184, 188 ff.; location of point of, 165; prevention of evaporation, 166, 168, 180; the pH change in the sample of, 166, 169; regulation of total CO_2 content, 166–167; pH control of, 167; addition of ammonia, 167, 169; effect of temperature on, 169, 171 ff., 183; precipitate, 169–171, 180 ff.; aeration method for study of, 165 ff., 173, 174; removal of biological activity from, 174–176; taking samples aseptically, 174 ff., 180 ff.; effect of modification of gas phase, 180; hydrogen ion concentration of, 182, 185; bacteriological count of, 182; Berkefeld controls, 186
Calcium solubility product constant, 147, 150, 151, 153, 155
California striped barnacle, 63. *See also* Marine organisms
California: tropical front development in, 323; interaction of tropical and polar maritime air between, and Hawaiian Islands, 327
Canadian oyster, sexual phases in, 120
Caprella scaura, 46
Caprellids, attachment of, to wood and cement, 42
Capsules, bacterial, 292
Carbohydrate fermentation, 291
Caribbean Sea, bottom samples taken in, 2
Cassidulina californica, 227
cushmani, 228
limbata, 227, 230
tortusa, 227, 228
Catalase: universal distribution of, 297; and metabolic activities, 297; and gen-

eral health and vigor, 297; determination of activity of, 298; found in different organs, 298, 304; in retractor muscles, 298, 304; from human blood, 299; from beef kidney, 299; mussel, 299, 300; relative activities of, from different tissues, 299, 300 (table 1); maximum activity of, 300, 305, 307; measurement of, 303; activity of, in young and mature animals, 303; optimum pH curves for, 305, 306; effect of temperature on, reaction, 307, 308, 309
Cellaria diffusa, 54
mandibulata, 54
Cellulose, digestion of, by bacteria, 291
Cephalization: marked in Nereis, 222; lack of, in Dendrostoma, 222
Cerebral ganglion, destruction of, 220–221
Chaetopterus, 52
Chamberlin, R. V., cited on sipunculids, 201
China Sea, origin of frontal systems in the, 321
Chromatin granules, 367
Chromidea, 365, 370
Chromidial: net, 365; theories, 366
Chromogenesis of marine bacteria, 291
Chromomeres, 367
Chromosomes of Patellina: telophase, 367, 368, 370; haploid, 367, 373; during first postzygotic division, 368; diploid, 368; in final metagamic divisions, 370–371; polarization of developing, 372; number of, in gametes, 373
Clyde Sea, nematodes and mollusks in muds of, 214
Clytia, 51
CO_2-producing bacteria contained in deep mud, 152
Cocci, percentage of, recovered from sea water, 292
Coe, Wesley R., **Season of Attachment and Rate of Growth of Sedentary Marine Organisms at the Pier of the Scripps Institution of Oceanography, La Jolla, California**, 37–86; **Development of the Gonads and the Sequence of Sexual Phases in the California Oyster (Ostrea lurida)**, 119–144
Coefficient, activity, of bicarbonate and carbonate ions in sea water, 264 ff. *See also* Sea water
Colorimetric method applied to sea water complex, 149
Copepods, marine: daily migrations of, 33; change in phototaxis of, 33; important part played by light in movements of, 34; compared with swimming algae, 34; resistance of, against poisons, 34, 35
Coprolitic Mud, 5, 14; color of, 7; organic and mineral constituents of, 7, 8; criteria for recognition of, 7, 8; from mouth of Congo River, 7; found in Gulf of Guinea, 8; chemical analyses of, 8; mode of origin not definitely known, 8; from coast of Panama, 8
Corallina cuvier var. calliptera, 50
gracilis, 47, 50
Cornuspira involvens, 51
Corycaeus rostratus, 33
venustus, 33
Crisia, 42
franciscana, 47, 54
geniculata, 47 *passim*, 54
pacifica, 47, 54
Crisulipora occidentalis, 54
Crucibulum spinosum, 63
Cummings, N. W., **A Sinker Method for the Determination of Specific Gravities**, 101–118
Cyclones: origin of, 321; development of secondary, 327
Cypselurus, 89
californicus, 89–93
Cytology of foraminifera: discussion, 365; description of gametes, 366; fertilization of zygotic nucleus, 367–368; pycnotic nuclei of degenerated gametes, 368; first postzygotic division, 368; origin of microspheric proloculum, 369; second postzygotic division, 369; final metagamic divisions, 370–371; escape of microspheric somatella, 371; development of megalospheric test, 371; syzygy and gametogenesis, 372; progamic divisions, 372; escape and multiple fission of somatella, 372; reduction division and gamete formation, 373

Darwin, cited on rôle of earthworm in alteration of soil, 210
Deep-sea bottom samples: area in which, were taken, 2–3; method employed in making analyses, 3, 12; symbols used to designate color, 5; determination of colloidal content, 19; chemical composition of, 20; occurrences of, 27; classification of, 28; environmental conditions of, 28

Dendrostoma, 201, 206, 210, 220
zostericola, 201–224
Dendrostoma zostericola: introvert of, 202, 206, 220, 221; tentacles, 202, 203, 204, 205, 208, 210, 218, 219; collar, 202, 203, 204, 211; nuchal organ, 203, 204, 220; mouth, 203; lips, 203; body cavity, 203; body fluid, 203, 204; cilia, 203; viscera, 203; pharynx, 203, 204; red blood corpuscles, 203; circulation of body fluid, 203; caeca, 203, 204; aeration of blood, 203; retractor muscles, 203, 204; ventral nerve cord, 204; circular muscles, 204; Polian vessel, 204; location of heart, 204; oesophagus, 204; nephridia, 204; digestive system, 204, 209; nervous system, 204; reproductive system, 204, 205, 206; positive thigmotropism of, 210; adaptation of, to stagnation of media, 211, 212; viability of, 211; response of, to oxygen, 212, 213, and to CO_2, 212; deprived of food and air, 212; laboratory experiments with, 212–213, 215–216; function of chemoreceptors, 112; production of hydrogen sulphide by, 213; removal from natural environment, 213; immersion in petroleum oil, 213; kept in moist chamber, 213; desiccation tolerance of, 213, 214; imbibition of dead organisms, 214; oxygen deprivation tolerance, 214, 215; methylene blue as intravitam stain, 215; effect of injury upon, 219; removal of tentacles, 219; narcotized with alcohol, 219, 220; limited regeneration of lost parts, 219, 221; respiration through skin, 219; destruction of cerebral ganglion, 220; integration of body parts, 220; effect of brain destruction on introvert, 220; lack of regeneration in brain, 221; severing of ventral nerve cord, 221; lack of cephalization, 222
Descriptions of Deep-Sea Bottom Samples from the Western North Atlantic and the Caribbean Sea, 1–31
Development of the Gonads and the Sequence of the Sexual Phases in the California Oyster (Ostrea lurida), 119–144
Diatoms: attachment of, to wood and cement, 42; many species of, found, 46; significance of metabolic activities of, 47; periodicity in abundance of, 50
Didemnum carnulentum, 48, 71
Discorbis, 370, 396, 402, 403
isabelleana, 51
Distaplia occidentalis, 48, 71
Dutch Harbor, air observations taken at, 324

Ectocarpus, 42 *passim*, 46, 49, 52
confervoides, 46, 49
cylindricus, 46, 49
granulosus, 46, 49
mitchellae, 49
mitchelli, 46
Egregia laevigata, 50
Electrolytes, supplied to mollusks by ocean environment, 307
Electrolytic dissociation, modern conception of, 235
Elphidium, 229, 402
articulatum, 227
hannai, 227, 229
hughesi, 227
spinatum, 227
Eponides ornata, 227
repanda, 227
Eucratea chelata, 54
Euherdmannia claviformis, 71
Eupomatus, 46
gracilis, 52
Euterpe acutifrons, 33, 34
Exocoetus, description of eggs of, 93

Fairchild, J. C., chemical analyses of deep-sea bottom samples made by, 4 *passim*
Feltham, Catharine B., and ZoBell, Claude E., **Preliminary Studies on the Distribution and Characteristics of Marine Bacteria**, 279–296
Fertilization cyst, 359, 360, 362, 372; formation of, 362; formation of zygotes within, 362; spherical zygotes within, 363; rupture of, 363; depressed nature of, 365; binucleate zygotic amoebulae within, 369; escape of microspheric agamonts from, 399
Fertilization nucleus, 402–403
Fisher, W. K., specimens of Dendrostoma identified by, 201
Flagella, bacterial, 292
Florida reef area, bacterial aspect of calcareous muds from, 146
Florkin, M. Marcel, cited on: coelmic fluid of Sipunculus nudus, 214; low oxygen content tolerance of sipunculids, 214; and their ability to store oxygen, 219

Föhn-heated air, 316, 322, 327, 329, 330, 332
Foraminifera: five faunal zones of, 227–228, 229; rate of sedimentation controls abundance of, 229; distribution and significance of, 355; growth cysts of, 358; pseudopodia of, 360; cytology of, 365 ff.; reorganization of protoplasm in, 366; dimorphism in, 402–403; nuclei of, 403
Fox, C. J. J., contributions of, to knowledge of factors involved in the buffer mechanism of sea water, 233
Fox, Denis L., and Marks, Graham W., **Studies on Catalase from the California Mussel,** 297–310
Fox, Denis L., and Peebles, Florence, **The Structure, Functions, and General Reactions of the Marine Sipunculid Worm, Dendrostoma zostericola,** 201–224
Free air soundings, importance of, 312
Frontal movements of air masses, 323 ff.
Frontal systems: movements of, of air masses, 323 ff.; of February 6 to March 6, 1930, 324 ff.; rate of movement of, across ocean, 325; turning of the occluded part of, 325
Frontogenesis, regions of, 320, 321, 322, 323
Frontolysis, zones of, 321

Galtsoff, P. S., cited on Ostrea virginica and O. gigas, 121
Gametes: protoplasm of, 362; size and movements of, 365, 367; description of, 366; fusion of, 366–367; nucleus of mature, 367; pycnotic nuclei of degenerated, 368; formation of, 373; determination of chromosome number in, 373; amoeboid, 373
Gametocytes: plasmotomy of, 362; mononucleate, 365
Gametogenesis, 362, 372, 373, 403; in Ostrea lurida, 124
Gee, Haldane, **Calcium Equilibrium in Sea Water. I. Theoretical Considerations,** 145–157; **IV. Apparatus for Eliminating Bacterial Activity in Chemical Studies,** 174–179; **V. Preliminary Experiments on Precipitation by Removal of Carbon Dioxide under Aseptic Conditions,** 180–187; **Bacteriological Water Sampler,** 191–200
Gee, Haldane, and Moberg, Erik G., **Calcium Equilibrium in Sea Water. III. Empirical Variation of Gas Phase,** 165–173
Gee, Haldane, Greenberg, D. M., and Moberg, Erik G., **Calcium Equilibrium in Sea Water. II. Equilibration in Sealed Bottles Shaken at Constant Temperature,** 158–164
Genes, transmission of hereditary characteristics by, 366
Gerould, J. H., cited on photic organs of Phascolosoma, 220
Gigantina spinosa, 50
Globigerina conglomerata, 51
Globigerina Ooze, 5, 9, 15; color of, 10; organic constituents of, 10; mineral particles found in, 10; dried, hard to crush, 10–11; difference between, and Red Clay, 11; distribution of, 11; chemical analyses of, 19, 20; occurrence of, 27
Globobulimina pacifica, 228
Grantia, 48, 51
Green Mud, 5, 14; a variety of Blue Mud, 6; color of, 6, 7
Greenberg, David M. *et al.*, **The Buffer Mechanism of Sea Water,** 231–278
Greenberg, D. M., Moberg, Erik G., and Gee, Haldane, **Calcium Equilibrium in Sea Water. II. Equilibration in Sealed Bottles Shaken at Constant Temperature,** 158–164
Growth cyst: in foraminifera, 358–359; of Patellina, 399
Gyropleurodus, 87
  francisci, 87

Hall Canyon, California: marine sediments exposed in, 225, 226, 229; fossil species collected in 226–227, 229–230
Hamberg, A., cited on carbon dioxide of sea water, 233
Hamburg: polar maritime air outbreak at, 320; comparison of air mass at Seattle with that at, 320
Hanzawa, S., statement of, concerning minute calcareous remains, 9
Hawaiian Islands: upper air data at, 315; transitional polar maritime air at, 318; Kona storms of, 322, 323; interaction of tropical and polar maritime air between California and, 327
Hemerythrine, 203, 214, 219

Hemitripterus marmoratus, 98
Hermaphroditism in the genus Ostrea, 136
Herposiphonia, 49
verticellata, 46, 50
Heterostichus, 98
rostratus, 98–99
Hilton, W. A., cited on nervous system of Phascolosoma, 204
Hippothoa, 47
hyalina, 54
Honolulu: free air soundings made at, 311; air observations made at, 324
Human blood: temperatures at which maximum catalase was obtained from, 307, 309; catalase thermal optima of, 309
Hydroclathrus clathrus, 50
Hydrogen ion concentration: in sea water, 235; method of determining, 248–249, 306; no arrangement in mollusks for control of, 307
Hydroids, seasons of maximum abundance of, 51

Icelandic cyclone of the Atlantic, 323
Interaction: of air masses, 320 ff.; of polar and maritime air, 327
Irving, L., cited on solubility of calcium in sea water, 147, 148, 151
Isogametes: spherical amoeboid, 362; zygote formed by two haploid, 366

Jackson, C. M., cited on Lumbricus terrestris, 312
Japan: origin of frontal systems in, 321; current, 321; origin of average North Pacific storm in, 323
Japanese Imperial Marine Observatory, daily meteorological charts of, for the North Pacific Ocean, analyzed, 311, 324
Java, upper air data at Batavia, 315
Johnston, J., cited on solubility of calcium carbonate, 147, 148, 149, 150

Kodiak, air observations made at, 324
Kona storms, Hawaiian Islands, development of, 322, 323
Krogh, A., measurement of carbon dioxide tension of sea water introduced by, 233
Kuro-Siwo. *See* Japan current

Leathesea, 42, 49
difformis, 46, 50
Legendre, R., cited on equilibrium relationships in sea water, 231
Lepeschkin, W. W., **Some Experiments on the Influence of Light and Poisons on Marine Copepods, with Reference to Their Daily Migration**, 33–35
Lichenopora, 42
radiata, 54
Life History of Patellina corrugata Williamson, a Foraminifer, The, 355–392
Literature cited on: Deep-Sea Bottom Samples, Atlantic and Caribbean, 29; Growth of Sedentary Marine Organisms, 74; Specific Gravities by Sinker Method, 118; Sexual Phases in the Oyster, 139; Calcium Equilibrium in Sea Water, 156, 164, 173, 179, 187, 190; Bacteriological Water Sampler, 200; Buffer Mechanism of Sea Water, 274; Marine Bacteria, 294; Catalase from California Mussel, 310; Air Masses of the North Pacific, 335, 336; Life History of Patellina corrugata Wil., 375; Further Studies on Patellina corrugata Wil., 404
Long Beach, California: bathymetric and temperature ranges of Recent foraminifera off, 225, 226–227, 229; material collected between Catalina Island and, 225
Los Angeles: air soundings made at, 311, 330; wind velocities observed in, Basin, 330
Low pressure: center, 322, 324; semipermanent, region of the Aleutian Islands, 323; latitudinal extension of the, trough, 326
Lumbricus terrestris, 213

McClendon, J. F., cited on calcium carbonate precipitation in sea water, 165
Manila, surface temperature of tropical current at, 324
Marine bacteria. *See* Bacteria
Marine organisms: economic problem of, 38; observations to be made on, 38, 39; findings concerning, 40, 41; free-swimming stages of, 41, 63; cyclical periodicity of reproductive period of, 41–42, 48; wooden and cement surfaces for attachment of, compared, 42, 44–45; sequence of appearance of, 46–49; sexual maturity of, 47; two seasons of maximum abundance of hydroids, 51; time of appearance and habits of annelids, 52; bryozoa, 53; larvae of mollusks,

54; reproductive period of Pacific coast oyster, 54–60; broad-eared scallop, 60–61; edible mussels found on blocks, 61–62; California striped barnacle, 63–70 *passim;* other crustacea, 70–71; ascidians, 71

Marine sediments, effect of chemical and physical alteration of, on worms, 210

Maritime histories of air masses, differences in, 320

Marks, Graham W., and Fox, Denis L., **Studies on Catalase from the California Mussel,** 297–310

Mass law equations: constants of, 249 ff.; evaluation of pK_1' and pK_2' of, 249 ff.; relation between pK_2' and pH in, 255 ff.; relation between pK_2' and salinity in, 263. *See also* Buffer mechanism; Sea water

Megalospheric agamont, 355; development of, 359; syzygy, 360–362; formation of ectoplasm, 360–361; no fusion of endoplasm, 361, 362; fertilization cyst, 362, 363; gametogenesis, 362; plasmotomy of gametocytes, 362; association of, in syzygy, 363, 401; migration of single nucleus of, 363; activities of cytoplasm, 363; somatella of, 363–365; haploid isogametes of, forming zygote, 366; escape of, from multiple fission cyst, 397, and from fertilization cyst, 399; rate of reproduction of, 401; variations in size and shape, 401; average number of young produced, 401; multinucleate, 403

Megalospheric generation, 355, 356

Megalospheric tests, 355; high proportion of microspheric tests to, 359, 401–402; development of, 360, 363, 371–372, 397; migration of single nucleus of, 362; differentiation of, 397; spiral chambers of, 397 *passim,* 398; line of pseudopodial pores, 397; aperture of, 397, 398; secondary septa of, 397; earliest free stage of, 397; proloculum of, 397, 398; terminal septum of, 397; crescent-shaped chamber of, 397–398; dimorphic nature of, 398; chamber forming continuous spire, 398; measurements of, 398, 399; addition of new chambers to, 399. *See also* Proloculum

Membranipora, 47
membranacea, 47, 54
tehuelcha, 54

Menipea occidentalis, 47, 54

Meyer, A., cited on: body fluid in Phalascosoma, 203; blood circulation through tentacles, 203

Microspheric agamont, 355; multiple fission cyst of, 359, 363; fertilization of, 359; rest period of, 360; development of, 363; liberation of, 363; four-celled somatella of, 369; rate of reproduction of, 399; size of initial nucleus of, 403. *See also* Proloculum; Somatella

Microspheric tests: high proportion of, to megalospheric tests, 359, 401–402; development of, 359–360, 363; escape of, from fertilization cyst, 360; growth of, 360; proloculum of, 396; spiral chamber of, 396; marginal chamberlets of, 396; earliest free stage of, 396; spiral stage of, 396; aperture of, 396, 397; discontinuous growth of, 396–397; median septa of adult, 397; semilunar chambers of, 397; dimorphic nature of, 398; elliptical base of, 398; measurements of, 398, 399; addition of new chambers to, 399; ratio of, to megalospheric cultures, 401

Midway Islands, air observations made at, 324

Moberg, Erik G., and Gee, Haldane, **Calcium Equilibrium in Sea Water. III. Empirical Variation of Gas Phase,** 165–173

Moberg, Erik G., Gee, Haldane, and Greenberg, D. M., **Calcium Equilibrium in Sea Water. II. Equilibration in Sealed Bottles Shaken at Constant Temperature,** 158–164

Moberg, Erik G., *et al.*, **The Buffer Mechanism of Sea Water,** 231–278.

Molgula verrucifera, 48, 71

Mollusks: nudibranch, 46, 62; Pacific coast oyster, 54–60; broad-eared scallop, 60–61; edible mussels found on blocks, 60–61; other forms of, 62–63; possess no control over salt concentration, 307; colloidal stability in body fluids of, 307; without control over body temperatures, 307; ocean environment supplies all needs of, 307. For catalase in mussels, *see* Catalase

Monergids, 365

Monsoon, winter of the far east, significance of, 316

Moore, H. B., cited on nematodes and mollusks in muds of Clyde Sea, 214

Morphogenesis of the Test and the Biological Significance of Dimorphism in the Foraminifer Patellina corrugata Williamson, 393–404
Mountain ranges, effect of, on air masses, 329, 330
Multiple fission, 360, 362, 365, 397; cyst, 359, 360, 363, 371, 397
Murray, John, work of, on sea bottom deposits, 4
Mussel catalase. *See* Catalase
Myers, Earl H., **The Life History of Patellina corrugata Williamson, a Foraminifer**, 355–392; **Morphogenesis of the Test and the Biological Significance of Dimorphism in the Foraminifer Patellina corrugata Williamson**, 393–404
Mytilus californianus, 61, 297
edulis, 61

Nagasaki, temperature of polar air current at, 324
Naha, temperature of tropical air at, 324
Natland, Manley L., **The Temperature- and Depth-Distribution of Some Recent Fossil Foraminifera in the Southern California Region**, 225–230
Nematodes, oxygen deprivation tolerance of, 214
Nereis, 222
agassizii, 46, 52
latescens, 52
vexillossa, 52
Nitrites: bacterial reduction of, 290; oxidation of, 291
Nonion umbilicatula, 228, 230
var. pacifica, 228
scapha, 227
North Atlantic, bottom samples taken in the, 2
North Pacific: synoptic charts of, 311; warmest part of, 321; coldest water of, 321; weather charts of, 324
North Pacific air masses, 311; modification of, 312; distribution of centers of interaction of, 321
Norton, R. D., cited on foraminiferal ecology, 3–4, 11
Notes on the Habits, Eggs, and Young of Some Fishes of Southern California, 87–99
Nucula tenuis, oxygen deprivation tolerance of, 214

Obelia, 42 *passim*, 49, 51
dichotoma, 46 *passim*, 47, 51
Ocean: needs of mollusks supplied by, 307; difference in surface temperature of, 321
Oithona brevicornis, 33
nana, 33
similis, 33
Oncaea venusta, 33, 34
Orient, difficulty of obtaining meteorological data from, 312
Orthopyxis caliculata, 51
everta, 51
Orton, J. H., cited on sexual phases of Ostrea edulis, 120, 125, 136
Ostrea edulis, 120, 124
equestris, 121
gigas, 121, 136
virginica, 41, 121
Ostrea lurida: sequence of sexual phases in, 120, 124, 125, 128, 129, 136–137; length of spawning season of, 120; discharge of ripe sexual products, 120–121, 128, 129; eggs of, fertilized within mantle cavity, 121; free-swimming stage of young, 121; reproductive system of, 121–124, 125, 132–133; sperm balls of, 125, 129, 131, 132; fertilization of ova, 125, 129, 131; generations of, in one year, 129; age of, at first spawning, 130; true hermaphroditism in, 130; typical protandry in, 131; spermatogenesis in, 131; second male phase in, 133; resting period of, 133; second female phase in, 134; later sexual phases, 134; wintering condition of, 134–135
Ovogomia, 122, 131
Oxygen deprivation: tolerance, 214; capacity to withstand, 215
Oxyhemerythrine, 214
Oxyjulus californicus, 98
Oyster, Pacific coast. *See* Ostrea lurida; Marine organisms

Pacific anticyclone, 322
Pacific Coast: meteorology of, 328 ff.; physiographic nature of, 328; free air soundings made on, 332; storm area of, 333
Pacific Ocean, continental air over, 329
Palitzsch, S., alkalinity and hydrogen ion concentration in sea water reviewed by, 233
Paracalanus parvus, 33, 34

Patellina, 396, 399, 402
protea, 393
subcretacea, 383
Patellina corrugata, 51, 356, 357, 366, 373–374; ecology of, 358; observations on, 358; rôle of cysts in life cycle of, 358–359; sexual and asexual generations in, 359; microspheric generation in, 359; fertilization of, 359; nuclei in adult, 370; region occupied by spindle in, 373; dimorphic tests of, 393, 396, 402–403; systematic position of, 393; measurement of tests of, 398, 399; growth cyst of, 399; rate of reproduction of, 399–402; mitotic division of, 403. *See also* Megalospheric agamont; Megalospheric tests; Microspheric agamonts; Microspheric tests.
Pearl Harbor: temperature and moisture content of air at, 315; tropical air curve at, 314
Pecten, 42
caurimus, 230
circularis var. aequisculatus, 63
latiauritus, 46, 60
Peebles, Florence, and Fox, Denis L., **The Structure, Functions, and General Reactions of the Marine Sipunculid Worm, Dendrostoma zostericola,** 201–224
pH curves, optimum: for catalase, 305, 306; suitable for successful living of mollusks, 307
Phalascosoma, 203, 204, 220
gouldii, 206
vulgare, 220
Photic organs, 220
Phototactic sensitiveness of marine copepods, 33–34
Phytomonads, alternation of generations in, 355
Pigmentation of marine bacteria, 291
Planulina ornata, 51
Plectofrondicularia californica, 228, 230
Plumularia setacea, 46, 51
Polar continental air masses, 312; stability of, 314; feeding of, into tropical regions, 316; föhn-heated, 316, 329; two types of, 329. *See also* Air masses
Polar maritime air masses, 312, 325; properties of, 314–315; contain less moisture, 318; comparison of, with European data, 320; interaction of tropical and, 327. *See also* Air masses
Polymorphine charlottensis, 227
"polynergid," 365
Polystomella, 355
crippa, 355, 373, 402, 403
Pourtalès, L. F., work of, on sea bottom deposits, 4
Preliminary Studies on the Distribution and Characteristics of Marine Bacteria, 279–296
Proloculum: microspheric, 359, 363, 369, 396, 402; megalospheric, 372, 397, 398, 402
Protozoa: association of, with other marine organisms, 51; rejection of chromidial theory for, 366; need for study of life cycle of, 366
Pteropod Ooze, 5, 9, 15; color of, 10; organic and mineral constituents of, 10; classification of, 10; distribution of, 10
Pullenia bulloides, 228 *passim*, 230
Pulvinulinella pacifica, 228

Quinqueloculina akneriana, 227, 402, 403
lamarckiana, 51

Rainfall: cause of, 316; on Pacific Coast, 333
Red Clays, 5; usual color of, 11; remains of organisms in, 11; basis for separation of, from Globigerina Ooze, 12; very coherent, 12; silt-clay percentages of, 12; comparison of, with Blue Mud, 15; dried, 16; chemical analyses of, 19; occurrence of, 27
"Repetto" beds of Ventura and Los Angeles basins, 226
Revelle, Roger, **Calcium Equilibrium in Sea Water. VI. Report on Precipitate Obtained by Removal of Carbon Dioxide from Sea Water,** 188–190
Revelle, Roger, *et al.*, **The Buffer Mechanism of Sea Water,** 231–278
Robertina charlottensis, 227
Robulus, 402
Rotalia beccarii, 227, 228 *passim*
Rotaliidae, 396
Roughly, T. C., cited on sexual phases in the Ostrea species, 136
Ruppin, E., cited on Ca and CO_2 in the water of the North Sea, 151

Sampling technic: for marine bacteria, 191, 279; for marine muds, 288

San Diego: free air soundings made at, 311, 316, 325, 328, 332; upper air data at, 312; tropical air curve at, 316; air inversions at, 319; quasistationary cold front in neighborhood of, 323; equivalent-potential temperature at, 325, 326, 327, 328, 334; tendency toward stratification noted at, 326; on edge of tropical current, 328, 333
Sand: ingested by Dendrostoma zostericola, 204, 209; used to stimulate responses, 208; nutrient matter obtained from, 209, 210
Santa Catalina Island: fossil material collected in channel between Long Beach and, 225; composition of southern portion of, 228; foraminifera found in sediment off coast of, 229
Santa Cruz Island, 225
Sarda chilensis, 87
Saxicava pholadis, 62
Scallops, broad-eared, 60–61. *See also* Marine organisms
Scorpaena, 93
guttata, 93, 94 *passim*
Scorpaenichthys, 98
marmoratus, 98
Scrupocellaria diegensis, 47, 54
Scytosiphon, 42
lomentaria, 46, 50
Sea water: concentration of hydrogen and hydroxyl ions in, 231, 233, 306; buffer mechanism in, 231–232; mass action law applied to, 233, 234; (complexity of, 234, 249 ff., 255, 263;) quantitative formulation of the buffer mechanism of, 235; variation of CO_2 in, 247–248; total pH of, 248–249; activity coefficients of, 264 ff.; CO_2 components and buffer capacity of, 268 ff.; calculation of CO_2 components of, 268, 270; relation of buffer capacity to pH of, 271; solubility of calcium carbonate in, 271; activity coefficient of calcium ion in, 272; fungi in, 292; molds in, 293. *See also* Buffer mechanism; Mass law equations
Season of Attachment and Rate of Growth of Sedentary Marine Organisms at the Pier of the Scripps Institution of Oceanography, La Jolla, California, 37–86
Seattle: free air soundings made at, 311, 317, 325, 330, 332, 334; upper air data at, 312; transitional polar maritime air at, 318–319; comparison of polar maritime data of, with European, 320; equivalent-potential temperature at, 325, 326, 327, 328; high moisture content of air at, 330; air overrunning at, 334
Serpula, 46
columbiana, 52
Sigmomorphina frondiculariformis, 227
Sinker Method for the Determination of Specific Gravities, A, 101–118
Sipunculid worms, 201; habitat of, 201, 204; uses of introvert made by, 204; low oxygen-content tolerance of, 214
Sipunculus nudus, coelomic fluid of, 214
Smittia collifera, 54
reticulata, 54
Somatella: nuclei of, 360 *passim*, 361, 363–364, 365; multiple fission of the microspheric, 360, 362; four-celled, of microspheric agamont, 369, 401; escape of microspheric, 371, 397; multinucleate, 397
Some Experiments on the Influence of Light and Poisons on Marine Copepods, with Reference to Their Daily Migrations, 33–35
Spärck, R., cited on sexual phases in Ostrea edulis, 124
Specific gravities: accurate determination of, 101; method for rapid determination of, 101, 102; theory of sinker method for determining, 103 ff.; computations for single determination of, 117
Spermatids, 125
Spermatocytes, 124, 125
Spermatogenesis: in Ostrea lurida, 124, 131–132; condition of genital ducts at completion of, 124; beginning of, 128; successive stages of, 129
Spermatogonia, 122, 124, 129, 131
Spermatophores, 131
Sphyraena argentea, 87
Spirillina, 370, 395, 396, 402
vivipara, 393
Spirorbis, 46, 52
Sporozoa, alternation of generations in, 355
Stafford, J., cited on sexual phases of the Canadian oyster, 120
Strait of Juan de Fuca, seaward flow of air through, 329
Structure, Functions, and General Re-

actions of the Marine Sipunculid Worm, Dendrostoma zostericola, The, 201–224
Studies on Catalase from the California Mussel, 297–310
Styela montereyensis, 71
Synidotea harfordi, 71
Syzygium, 359, 360, 362, 401
Syzygy, 360, 362, 401; and gametogenesis, 372

Temperature: effect of, on marine bacteria, 282; storage, of water samples, 283; influence of, on chromogenesis, 291; effect of, on catalase reaction, 307; differences in ocean surface, 321; contrast in, in two air currents, 324; high equivalent-potential, at San Diego and Seattle, 325, 327, 328
Temperature-activity curve for mussel catalase, 308
Temperature- and Depth-Distribution of Some Recent and Fossil Foraminifera in the Southern California Region, The, 225–230
Temperature, body: of mollusks, 307; capacity of homoiothermic forms to regulate, 307
Temperature diagram, equivalent-potential, 313
Tentacles, Dendrostoma zostericola: used for obtaining food, 208, for interchange of gases, 208, and for breathing, 210, 212; reaction of, to CO_2, 216, to the action of salt solution, 217, and to light, 218; results of amputation of, 219, 221
Thalmoporella, 47
rozieri, 47
californiensis, 54
Thorp, E. M., **Descriptions of Deep-Sea Bottom Samples from the Western North Atlantic and the Caribbean Sea,** 1–31
Transitional polar continental air mass: effect of winter monsoon on cold type of, 316; occurrences of, 316; warm type of, 316; stability of, 316; high moisture content of, 317, 318. *See also* Air masses
Transitional polar maritime air mass: moist type of, 318–319; dry type of, 319; föhn as it occurs in, 329
Transitional tropical maritime air masses, 319; characterized by widespread fog conditions, 319–320
Tretomphalus bulloides, 51
Triloculina circularis, 51
trigonula, 227
Tropical maritime air masses, 312, 315; properties of, 315; change caused by northward movement of, 315–316; interaction of polar and, 327. *See also* Air masses.

United States: continental shelf of, 2; frontal developments along coast of, 323; types of frontogenesis in, 323
United States Bureau of Soils, analyses of deep-sea bottom samples made by, 2, 4, 19
United States Coast and Geodetic Survey, collection of deep-sea bottom samples made by, 2, 4
United States Geological Survey, analyses of deep-sea bottom samples made by, 3
United States Weather Bureau, San Francisco, charts of, analyzed, 311, 324
Urolophus halleri, 88, 89
Uvigerina peregrina, 228, 230

Van Slyke's manometric apparatus, 159, 181
Vaughan, T. W., cited on relationship of other sciences to Oceanography, 145–146
Ventura anticline, 226
Vermetids, 63

Weisensee, H., cited on sex conditions in Anodonta, 130
Wilson, Charles B., identification of copepods by, 33
Winter monsoon. *See* Monsoon
Worms. *See under* various generic names

Yangtse Valley, frontogenesis in, 321
Yukon Valley: wind currents of, 322; air of, 323

ZoBell, Claude E., and Feltham, Catharine B., **Preliminary Studies on the Distribution and Characteristics of Marine Bacteria,** 279–296
Zoöbotryon pellucida, 54
Zoöspores, flagellate, 365, 367
Zygotes: origin of, 355, 359; formation of, 362, 363, 371, 401; held together by hyaline cytoplasm, 363; formation of larger, 365
Zygotic: amoebula, 359, 369; nucleus, 359, 363, 367

CONTENTS

PAGE

1. Descriptions of Deep-Sea Bottom Samples from the Western North Atlantic and the Caribbean Sea, by E. M. Thorp 1

2. Some Experiments on the Influence of Light and Poisons on Marine Copepods, with Reference to Their Daily Migration, by W. W. Lepeschkin 33

3. Season of Attachment and Rate of Growth of Sedentary Marine Organisms at the Pier of the Scripps Institution of Oceanography, La Jolla, California, by Wesley R. Coe 37

4. Notes on the Habits, Eggs, and Young of Some Fishes of Southern California, by Percy Spencer Barnhart 87

5. A Sinker Method for Determination of Specific Gravities, by N. W. Cummings ... 101

6. Development of the Gonads and the Sequence of Sexual Phases in the California Oyster (*Ostrea lurida*), by Wesley R. Coe 119

7. Calcium Equilibrium in Sea Water. I. Theoretical Considerations. II. Equilibration in Sealed Bottles Shaken at Constant Temperatures. III. Empirical Variation of Gas Phase. IV. Apparatus for Eliminating Bacterial Activity in Chemical Studies. V. Preliminary Experiments on Precipitation by Removal of Carbon Dioxide under Aseptic Conditions. VI. Report on Precipitate Obtained by Removal of Carbon Dioxide from Sea Water, by Haldane Gee in collaboration with Erik G. Moberg, D. M. Greenberg, and Roger Revelle 145

8. Bacteriological Water Sampler, by Haldane Gee 191

9. The Structure, Functions, and General Reactions of the Marine Sipunculid Worm, *Dendrostoma zostericola*, by Florence Peebles and Denis L. Fox 201

10. The Temperature- and Depth-Distribution of Some Recent and Fossil Foraminifera in the Southern California Region, by Manley L. Natland 225

11. The Buffer Mechanism of Sea Water, by Erik G. Moberg, David M. Greenberg, Roger Revelle, and Esther C. Allen 231

12. Preliminary Studies on the Distribution and Characteristics of Marine Bacteria, by Claude E. ZoBell and Catharine B. Feltham 279

13. Studies on Catalase from the California Mussel, by Graham W. Marks and Denis L. Fox 297

14. The Air Masses of the North Pacific, by Horace Robert Byers 311

15. The Life History of *Patellina corrugata* Williamson, a Foraminifer, by Earl H. Myers 355

16. Morphogenesis of the Test and the Biological Significance of Dimorphism in the Foraminifer, *Patellina corrugata* Williamson, by Earl H. Myers 393

Index 405